


AMENDMENTS TO THE CLAIMS

1. (Currently amended) An aminoquinoxaline compound of the following formula (1a)

~~—[Chemical Formula 1]~~



wherein R<sup>1</sup> and R<sup>2</sup> independently represent a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>1</sup> and R<sup>2</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

R<sup>3</sup> and R<sup>4</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be

substituted with Y provided that when R<sup>3</sup> and R<sup>4</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

X<sup>1</sup> represents -NH-R<sup>5</sup>-NH<sub>2</sub> [[or -NH-R<sup>6</sup>]];

R<sup>5</sup> represents a ~~C<sub>4</sub>-C<sub>10</sub> alkylene group, a C(O)CH<sub>2</sub>, —CH<sub>2</sub>C(O), a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a pyrrole ring which may be substituted with Y, a furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;~~

~~—R<sup>6</sup> represents a hydrogen atom, a C<sub>4</sub>-C<sub>10</sub> alkyl group, an acetyl group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y;~~


Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or

a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group provided that if Z is two or more in number, Z may be the same or different.

2. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein R<sup>1</sup> and R<sup>2</sup> in the above formula (1) independently represent a group of the following formula (2)

~~—[Chemical Formula 2]~~




wherein R<sup>7</sup>–R<sup>11</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>4</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>4</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a

C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group.

3. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein R<sup>1</sup> and R<sup>2</sup> in the above formula (1) independently represent a group of the following formula (3)

~~—[Chemical Formula 3]~~




wherein R<sup>12</sup>–R<sup>18</sup> independently represent, each substituted at an arbitrary position on the ring of the formula, a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group.

4. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein R<sup>1</sup> and R<sup>2</sup> in the above formula (1) independently represent a group of the following formula (4)

~~—[Chemical Formula 4]~~




wherein  $R^{19}$ – $R^{21}$  independently represent, each substituted at an arbitrary position on the ring of the formula, a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  haloalkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a  $C_1$ – $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group; and

$A^1$  represents NH, O or S.

5. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein  $R^1$  and  $R^2$  in the above formula (1) represent a group of the following formula (5)

[Chemical Formula 5]




wherein  $R^{22}$  represents a halogen atom or a cyano group,  $R^{23}$ – $R^{26}$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  haloalkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a  $C_1$ – $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group.

6. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein  $R^5$  in the formula (1) represents a group of the following formula (6)

[Chemical Formula 6]




wherein  $R^{27}$ – $R^{30}$  independently represent, each substituted at an arbitrary position on the ring of the formula, a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrolyl group, a furyl group or a condensed heteroaryl group.

7. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein R<sup>5</sup> in the formula (1) represents a group of the following formula (7)

{Chemical Formula 7}




wherein R<sup>31</sup>-R<sup>32</sup> independently represent, each substituted at an arbitrary position on the ring of the formula, a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrolyl group, a furyl group or a condensed heteroaryl group; and

W<sup>1</sup> represents NH, O or S.

8. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein  $R^5$  in the formula (1) represents a group of the following formula (8)

[Chemical Formula-8]



wherein  $R^{33}$ – $R^{34}$  independently represent, each substituted at an arbitrary position on the ring of the formula, a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z;


Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  haloalkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a  $C_1$ – $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group; and

$Q^1$  represents NH, O or S.

9. **(Currently amended)** The aminoquinoxaline compound according to claim 1, wherein  $R^5$  in the formula (1) represents a group of the following formula (9)

~~[Chemical Formula 9]~~






wherein  $R^{35}$ – $R^{40}$  independently represent, each substituted at an arbitrary position on the ring of the formula, a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and


Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  haloalkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a  $C_1$ – $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group.

10. **(Cancelled)**

11. **(Currently amended)** ~~The aminoquinoxaline compound according to claim 1, wherein  $R^6$  in the formula (1) represents a group~~ An aminoquinoxaline compound of the following formula (11)

~~{Chemical Formula 11}~~





wherein R<sup>1</sup> and R<sup>2</sup> independently represent a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>1</sup> and R<sup>2</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

R<sup>3</sup> and R<sup>4</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>3</sup> and R<sup>4</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$R^{46}$ – $R^{48}$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z;



Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  haloalkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a  $C_1$ – $C_{10}$  cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ – $C_{10}$  alkyl group, a  $C_1$ – $C_{10}$  haloalkyl group, a  $C_1$ – $C_{10}$  alkoxy group, a  $C_1$ – $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group provided that if Z is two or more in number, Z may be the same or different; and

$W^2$  represents NH, O or S.

12. **(Currently amended)** ~~The aminoquinoxaline compound according to claim 1, wherein~~  
 ~~$R^6$  in the formula (1) represents a group~~ An aminoquinoxaline compound of the following  
formula (12)

[Chemical Formula 12]



wherein  $R^1$  and  $R^2$  independently represent a hydrogen atom, a hydroxyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when  $R^1$  and  $R^2$  are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$R^3$  and  $R^4$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a

furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>3</sup> and R<sup>4</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;



R<sup>49</sup>-R<sup>51</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group provided that if Z is two or more in number, Z may be the same or different; and

Q<sup>2</sup> represents NH, O or S.

13. **(Currently amended)** ~~The aminoquinoxaline compound according to claim 1, wherein~~  
~~R<sup>6</sup> in the formula (1) represents a group~~ An aminoquinoxaline compound of the following  
formula (13)

~~{Chemical Formula 13}~~



wherein R<sup>1</sup> and R<sup>2</sup> independently represent a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>1</sup> and R<sup>2</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

R<sup>3</sup> and R<sup>4</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>3</sup> and R<sup>4</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;



R<sup>52</sup>-R<sup>58</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a

C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrolyl group, a furyl group or a condensed heteroaryl group provided that if Z is two or more in number, Z may be the same or different.

14. **(Currently amended)** ~~The aminoquinoxaline compound according to claim 1, wherein the group formed by bonding R<sup>1</sup> and R<sup>2</sup> through a single bond in the formula (1) is represented by~~  
the An aminoquinoxaline compound of the following formula (14)

~~—[Chemical Formula 14]~~



wherein R<sup>3</sup> and R<sup>4</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group



which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>3</sup> and R<sup>4</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

X<sup>1</sup> represents -NH-R<sup>5</sup>-NH<sub>2</sub> or -NH-R<sup>6</sup>;

R<sup>5</sup> represents a C<sub>1</sub>-C<sub>10</sub> alkylene group, a -C(O)CH<sub>2</sub>-, -CH<sub>2</sub>C(O)-, a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a pyrrole ring which may be substituted with Y, a furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;

R<sup>6</sup> represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, an acetyl group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl

group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;


A<sub>2</sub> are each  $[[\text{CN}]] \text{C}$  or N,

R<sup>59</sup>–R<sup>66</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>–C<sub>10</sub> alkyl group, a C<sub>1</sub>–C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>–C<sub>10</sub> alkyl group, a C<sub>1</sub>–C<sub>10</sub> haloalkyl group, a C<sub>1</sub>–C<sub>10</sub> alkoxy group, a C<sub>1</sub>–C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group provided that if Z is two or more in number, Z may be the same or different, provided that when A<sup>2</sup> represents N, R<sup>62</sup> and R<sup>63</sup> are both non-existent.

15. **(Currently amended)** An aminoquinoxaline compound of the following formula (1b),

–[Chemical Formula 15]



(1b)

wherein R<sup>1'</sup> and R<sup>2'</sup> join together to form -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-,  
-OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>SCH<sub>2</sub>-,  
-CH<sub>2</sub>CH<sub>2</sub>N(R')-, -N(R')CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>N(R')CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>O-,  
-OCH<sub>2</sub>CH<sub>2</sub>O-, -SCH<sub>2</sub>CH<sub>2</sub>S-, -OCH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH=CH-,  
-CH=CHCH<sub>2</sub>-, -OCH=CH-, -CH=CHO-, -SCH=CH-, -CH=CHS-,  
-N(R')CH=CH-, -CH=CHN(R')-, -OCH=N-, -N=CHO-, -SCH=N-,  
-N=CHS-, -N(R')CH=N-, -N=CHN(R')-, -N(R')N=CH-, -CH=N(R')N-,  
-CH=CHCH=CH-, -OCH<sub>2</sub>CH=CH-, -CH=CHCH<sub>2</sub>O-, -N=CHCH=CH-,  
-CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-, or -CH=NCH=N- wherein a hydrogen atom  
bonded to a carbon atom of these groups may be substituted with Y, and R' represents a  
hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a  
phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z,  
a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted  
with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be  
substituted with Z, a furyl group which may be substituted with Z, or a condensed heteroaryl  
group which may be substituted with Z;

R<sup>3</sup> and R<sup>4</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a  
nitro group, an amino group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which  
may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group  
which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl  
group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a  
furyl group which may be substituted with Y or a condensed heteroaryl group which may be

substituted with Y provided that when  $R^3$  and  $R^4$  are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$X^1$  represents  $-NH-R^5-NH_2$  or  $-NH-R^6$ ;

$R^5$  represents a  $C_1$ - $C_{10}$  alkylene group,  $-C(O)CH_2-$ ,  $-CH_2C(O)-$ , a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a pyrrole ring which may be substituted with Y, a furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;


$R^6$  represents a hydrogen atom, a  $C_1$ - $C_{10}$  alkyl group, an acetyl group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or

a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group, provided that when Z are two or more in number, Z may be the same or different.

16. **(Currently amended)** An aminoquinoxaline compound of the following formula (1c),  
[Chemical Formula 16]



wherein R<sup>1</sup> and R<sup>2</sup> independently represent a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>1</sup> and R<sup>2</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$R^{3'}$  and  $R^{4'}$  join together to form  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{O}-$ ,  
 $-\text{OCH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{OCH}_2-$ ,  $-\text{OCH}_2\text{O}-$ ,  $-\text{CH}_2\text{CH}_2\text{S}-$ ,  $-\text{SCH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{SCH}_2-$ ,  
 $-\text{CH}_2\text{CH}_2\text{N}(\text{R}')-$ ,  $-\text{N}(\text{R}')\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{N}(\text{R}')\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{O}-$ ,  $-\text{OCH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{OCH}_2-$ ,  $-\text{CH}_2\text{OCH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{OCH}_2\text{O}-$ ,  
 $-\text{OCH}_2\text{CH}_2\text{O}-$ ,  $-\text{SCH}_2\text{CH}_2\text{S}-$ ,  $-\text{OCH}_2\text{CH}_2\text{S}-$ ,  $-\text{SCH}_2\text{CH}_2\text{O}-$ ,  $-\text{CH}_2\text{CH}=\text{CH}-$ ,  
 $-\text{CH}=\text{CHCH}_2-$ ,  $-\text{OCH}=\text{CH}-$ ,  $-\text{CH}=\text{CHO}-$ ,  $-\text{SCH}=\text{CH}-$ ,  $-\text{CH}=\text{CHS}-$ ,  
 $-\text{N}(\text{R}')\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CHN}(\text{R}')-$ ,  $-\text{OCH}=\text{N}-$ ,  $-\text{N}=\text{CHO}-$ ,  $-\text{SCH}=\text{N}-$ ,  
 $-\text{N}=\text{CHS}-$ ,  $-\text{N}(\text{R}')\text{CH}=\text{N}-$ ,  $-\text{N}=\text{CHN}(\text{R}')-$ ,  $-\text{N}(\text{R}')\text{N}=\text{CH}-$ ,  $-\text{CH}=\text{N}(\text{R}')\text{N}-$ ,  
 $-\text{CH}=\text{CHCH}=\text{CH}-$ ,  $-\text{OCH}_2\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CHCH}_2\text{O}-$ ,  $-\text{N}=\text{CHCH}=\text{CH}-$ ,  
 $-\text{CH}=\text{CHCH}=\text{N}-$ ,  $-\text{N}=\text{CHCH}=\text{N}-$ ,  $-\text{N}=\text{CHN}=\text{CH}-$ , or  $-\text{CH}=\text{NCH}=\text{N}-$  wherein a hydrogen atom  
bonded to a carbon atom of these groups may be substituted with Y, and  $\text{R}'$  represents a  
hydrogen atom, a  $\text{C}_1$ - $\text{C}_{10}$  alkyl group, a  $\text{C}_1$ - $\text{C}_{10}$  haloalkyl group, a  $\text{C}_1$ - $\text{C}_{10}$  cyanoalkyl group, a  
phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z,  
a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted  
with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be  
substituted with Z, a furyl group which may be substituted with Z, or a condensed heteroaryl  
group which may be substituted with Z;

$\text{X}^1$  represents  $-\text{NH}-\text{R}^5-\text{NH}_2$  or  $-\text{NH}-\text{R}^6$ ;

$\text{R}^5$  represents a  $\text{C}_1$ - $\text{C}_{10}$  alkylene group,  $-\text{C}(\text{O})\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(\text{O})-$ , a divalent benzene ring  
which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a  
divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which  
may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a pyrrole

ring which may be substituted with Y, a furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;


R<sup>6</sup> represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, an acetyl group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group, provided that when Z are two or more in number, Z may be the same or different.

17. **(Currently amended)** An aminoquinoxaline compound of the following formula (1d),

[Chemical Formula 17]



wherein R<sup>1'</sup> and R<sup>2'</sup> join together to form -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-,  
 -OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>SCH<sub>2</sub>-,  
 -CH<sub>2</sub>CH<sub>2</sub>N(R')-, -N(R')CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>N(R')CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>O-,  
 -OCH<sub>2</sub>CH<sub>2</sub>O-, -SCH<sub>2</sub>CH<sub>2</sub>S-, -OCH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH=CH-,  
 -CH=CHCH<sub>2</sub>-, -OCH=CH-, -CH=CHO-, -SCH=CH-, -CH=CHS-,  
 -N(R')CH=CH-, -CH=CHN(R')-, -OCH=N-, -N=CHO-, -SCH=N-,  
 -N=CHS-, -N(R')CH=N-, -N=CHN(R')-, -N(R')N=CH-, -CH=N(R')N-,  
 -CH=CHCH=CH-, -OCH<sub>2</sub>CH=CH-, -CH=CHCH<sub>2</sub>O-, -N=CHCH=CH-,  
 -CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-, or -CH=NCH=N- wherein a hydrogen atom  
 bonded to a carbon atom of these groups may be substituted with Y, and R' represents a  
 hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a  
 phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z,  
 a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted  
 with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be  
 substituted with Z, a furyl group which may be substituted with Z, or a condensed heteroaryl  
 group which may be substituted with Z, and R<sup>3'</sup> and R<sup>4'</sup> join together to form



-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>S-,  
-SCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>SCH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>N(R')-, -N(R')CH<sub>2</sub>CH<sub>2</sub>-,  
-CH<sub>2</sub>N(R')CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>O-, -SCH<sub>2</sub>CH<sub>2</sub>S-,  
-OCH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH=CH-, -CH=CHCH<sub>2</sub>-, -OCH=CH-,  
-CH=CHO-, -SCH=CH-, -CH=CHS-, -N(R')CH=CH-, -CH=CHN(R')-,  
-OCH=N-, -N=CHO-, -SCH=N-, -N=CHS-, -N(R')CH=N-, -N=CHN(R')-,  
-N(R')N=CH-, -CH=N(R')N-, -CH=CHCH=CH-, -OCH<sub>2</sub>CH=CH-,  
-CH=CHCH<sub>2</sub>O-, -N=CHCH=CH-, -CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-,

or -CH=NCH=N- wherein a hydrogen atom bonded to a carbon atom of these groups may be substituted with Y, and R' represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z;

X<sup>1</sup> represents -NH-R<sup>5</sup>-NH<sub>2</sub> or -NH-R<sup>6</sup>;

R<sup>5</sup> represents a C<sub>1</sub>-C<sub>10</sub> alkylene group, -C(O)CH<sub>2</sub>-, -CH<sub>2</sub>C(O)-, a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a pyrrole ring which may be substituted with Y, a furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;


R<sup>6</sup> represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, an acetyl group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group, provided that when Z are two or more in number, Z may be the same or different.

18. **(Currently amended)** The aminoquinoxaline compound according to claim 15, wherein the group formed by joining R<sup>1'</sup> and R<sup>2'</sup> together is of the following formula (15)

[~~Chemical Formula 18~~]




wherein  $A^3$  represents O or S, and  $R^{67}$ – $R^{70}$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group.

19. **(Currently amended)** The aminoquinoxaline compound according to claim 16, wherein the group formed by joining  $R^{3'}$  and  $R^{4'}$  together is of the following formula (16)

~~—[Chemical Formula 19]~~




wherein  $A^4$  represents O or S, and  $R^{71}$ – $R^{74}$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrolyl group, a furyl group or a condensed heteroaryl group.

20. **(Currently amended)** The aminoquinoxaline compound according to claim 16, wherein the group formed by joining R<sup>3'</sup> and R<sup>4'</sup> together is of the following formula (17)

~~—[Chemical Formula 20]~~




wherein R<sup>75</sup> and R<sup>76</sup> independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrolyl group, a furyl group or a condensed heteroaryl group.

21. **(Currently amended)** A polyaminoquinoxaline compound having recurring units of the following formula (18a) ~~obtained by polymerizing the monomer defined in claim 1,~~

[Chemical Formula 21]



wherein  $R^1$  and  $R^2$  independently represent a hydrogen atom, a hydroxyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when  $R^1$  and  $R^2$  are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$R^3$  and  $R^4$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when  $R^3$  and  $R^4$  are, respectively, the above-defined phenyl,

pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$X^2$  represents  $-NH-R^{77}-NH-$  or  $-NH-R^{78}-$ ;

$R^{77}$  and  $R^{78}$  independently represent a  $C_1$ - $C_{10}$  alkylene group, a  $-C(O)CH_2-$ ,  $-CH_2C(O)-$ , a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a divalent pyrrole ring which may be substituted with Y, a divalent furan ring which may be substituted with Y or a condensed hetero ring which may be substituted with Y;


Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group provided that if Z is two or more in number, Z may be the same or different; and

~~n is an integer of 2 or over~~ n is an integer of 2 or more.

22. **(Currently amended)** A polyaminoquinoxaline compound having recurring units of the following formula (18b) obtained by polymerizing the monomer defined in claim 15,

—[Chemical Formula 22]



wherein R¹' and R²' join together to form -CH₂CH₂CH₂-, -CH₂CH₂O-,  
 -OCH₂CH₂-, -CH₂OCH₂-, -OCH₂O-, -CH₂CH₂S-, -SCH₂CH₂-, -CH₂SCH₂-,  
 -CH₂CH₂N(R')-, -N(R')CH₂CH₂-, -CH₂N(R')CH₂-, -CH₂CH₂CH₂CH₂-,  
 -CH₂CH₂CH₂O-, -OCH₂CH₂CH₂-, -CH₂CH₂OCH₂-, -CH₂OCH₂CH₂-, -CH₂OCH₂O-,  
 -OCH₂CH₂O-, -SCH₂CH₂S-, -OCH₂CH₂S-, -SCH₂CH₂O-, -CH₂CH=CH-,  
 -CH=CHCH₂-, -OCH=CH-, -CH=CHO-, -SCH=CH-, -CH=CHS-,  
 -N(R')CH=CH-, -CH=CHN(R')-, -OCH=N-, -N=CHO-, -SCH=N-,  
 -N=CHS-, -N(R')CH=N-, -N=CHN(R')-, -N(R')N=CH-, -CH=N(R')N-,  
 -CH=CHCH=CH-, -OCH₂CH=CH-, -CH=CHCH₂O-, -N=CHCH=CH-,  
 -CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-, or -CH=NCH=N- wherein a hydrogen atom  
 bonded to a carbon atom of these groups may be substituted with Y, and R' represents a  
 hydrogen atom, a C₁-C₁₀ alkyl group, a C₁-C₁₀ haloalkyl group, a C₁-C₁₀ cyanoalkyl group, a  
 phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z,  
 a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted

with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z;

$R^3$  and  $R^4$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when  $R^3$  and  $R^4$  are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$X^2$  represents -NH- $R^{77}$ -NH- or -NH- $R^{78}$ -;

$R^{77}$  and  $R^{78}$  independently represent a  $C_1$ - $C_{10}$  alkylene group, -C(O)CH<sub>2</sub>-, -CH<sub>2</sub>C(O)-, a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a divalent pyrrole ring which may be substituted with Y, a divalent furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl




group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group, provided that when Z are two or more in number, Z may be the same or different; and

~~n is an integer of 2 or over~~ n is an integer of 2 or more.

23. **(Currently amended)** A polyaminoquinoxaline compound having recurring units of the following formula (18c) obtained by polymerizing the monomer defined in claim 16,

~~—[Chemical Formula 23]~~



wherein R<sup>1</sup> and R<sup>2</sup> independently represent a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl

group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when R<sup>1</sup> and R<sup>2</sup> are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

R<sup>3</sup> and R<sup>4</sup> join together to form -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-,  
-OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>SCH<sub>2</sub>-,  
-CH<sub>2</sub>CH<sub>2</sub>N(R')-, -N(R')CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>N(R')CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>O-,  
-OCH<sub>2</sub>CH<sub>2</sub>O-, -SCH<sub>2</sub>CH<sub>2</sub>S-, -OCH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH=CH-,  
-CH=CHCH<sub>2</sub>-, -OCH=CH-, -CH=CHO-, -SCH=CH-, -CH=CHS-,  
-N(R')CH=CH-, -CH=CHN(R')-, -OCH=N-, -N=CHO-, -SCH=N-,  
-N=CHS-, -N(R')CH=N-, -N=CHN(R')-, -N(R')N=CH-, -CH=N(R')N-,  
-CH=CHCH=CH-, -OCH<sub>2</sub>CH=CH-, -CH=CHCH<sub>2</sub>O-, -N=CHCH=CH-,  
-CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-, or -CH=NCH=N- wherein a hydrogen atom bonded to a carbon atom of these groups may be substituted with Y, and R' represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z, or a condensed heteroaryl group which may be substituted with Z;

X<sup>2</sup> represents -NH-R<sup>77</sup>-NH- or -NH-R<sup>78</sup>-;

$R^{77}$  and  $R^{78}$  independently represent a  $C_1$ - $C_{10}$  alkylene group,  $-C(O)CH_2-$ ,  $-CH_2C(O)-$ , a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a divalent pyrrole ring which may be substituted with Y, a divalent furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;


Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group, provided that when Z are two or more in number, Z may be the same or different; and

~~n is an integer of 2 or over~~ n is an integer of 2 or more.

24. **(Currently amended)** A polyaminoquinoxaline compound having recurring units of the following formula (18d) obtained by polymerizing the monomer defined in claim 17,

—[Chemical Formula 24]



wherein R<sup>1'</sup> and R<sup>2'</sup> join together to form -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-,  
 -OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>SCH<sub>2</sub>-,  
 -CH<sub>2</sub>CH<sub>2</sub>N(R')-, -N(R')CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>N(R')CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>O-,  
 -OCH<sub>2</sub>CH<sub>2</sub>O-, -SCH<sub>2</sub>CH<sub>2</sub>S-, -OCH<sub>2</sub>CH<sub>2</sub>S-, -SCH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH=CH-,  
 -CH=CHCH<sub>2</sub>-, -OCH=CH-, -CH=CHO-, -SCH=CH-, -CH=CHS-,  
 -N(R')CH=CH-, -CH=CHN(R')-, -OCH=N-, -N=CHO-, -SCH=N-,  
 -N=CHS-, -N(R')CH=N-, -N=CHN(R')-, -N(R')N=CH-, -CH=N(R')N-,  
 -CH=CHCH=CH-, -OCH<sub>2</sub>CH=CH-, -CH=CHCH<sub>2</sub>O-, -N=CHCH=CH-,  
 -CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-, or -CH=NCH=N- wherein a hydrogen atom  
 bonded to a carbon atom of these groups may be substituted with Y, and R' represents a  
 hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a  
 phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z,  
 a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted  
 with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be

substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z, and R<sup>3'</sup> and R<sup>4'</sup> join together to form

-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>S-,  
-SCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>SCH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>N(R')-, -N(R')CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>N(R')CH<sub>2</sub>-,  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>-,  
-CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>OCH<sub>2</sub>O-, -OCH<sub>2</sub>CH<sub>2</sub>O-, -SCH<sub>2</sub>CH<sub>2</sub>S-, -OCH<sub>2</sub>CH<sub>2</sub>S-,  
-SCH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH=CH-, -CH=CHCH<sub>2</sub>-, -OCH=CH-, -CH=CHO-,  
-SCH=CH-, -CH=CHS-, -N(R')CH=CH-, -CH=CHN(R')-, -OCH=N-,  
-N=CHO-, -SCH=N-, -N=CHS-, -N(R')CH=N-, -N=CHN(R')-,  
-N(R')N=CH-, -CH=N(R')N-, -CH=CHCH=CH-, -OCH<sub>2</sub>CH=CH-,  
-CH=CHCH<sub>2</sub>O-, -N=CHCH=CH-, -CH=CHCH=N-, -N=CHCH=N-, -N=CHN=CH-,

or -CH=NCH=N- wherein a hydrogen atom bonded to a carbon atom of these groups may be substituted with Y, and R' represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z;

X<sup>2</sup> represents -NH-R<sup>77</sup>-NH- or -NH-R<sup>78</sup>-;

R<sup>77</sup> and R<sup>78</sup> independently represent a C<sub>1</sub>-C<sub>10</sub> alkylene group, -C(O)CH<sub>2</sub>-, -CH<sub>2</sub>C(O)-, a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be

substituted with Y, a divalent pyrrole ring which may be substituted with Y, a divalent furan ring which may be substituted with Y or a condensed hetero ring which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different; and

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> haloalkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrrolyl group, a furyl group or a condensed heteroaryl group, provided that when Z are two or more in number, Z may be the same or different; and

~~n is an integer of 2 or over~~ n is an integer of 2 or more.

25. **(Currently Amended)** A film obtained by use of an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

26. (Original) The film according to claim 25, wherein the film is prepared by spin coating, casting or vacuum deposition.

27. (Original) The film according to claim 25, wherein the film is obtained by compression molding.

28. (**Currently amended**) An electro chromic device made by use of an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

29. (**Currently amended**) A semiconductor device made by use of an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

30. (**Currently amended**) A p-type semiconductor obtained by oxidizing, with an oxidizing agent or through electrochemical doping, an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

31. (**Currently amended**) An n-type semiconductor obtained by reducing, with a reducing agent or through electrochemical doping, an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

32. (**Currently amended**) A solar cell made by use of ~~the p-type semiconductor defined in claim 30 and the n-type semiconductor defined in claim 31~~ a p-type semiconductor and an n-type semiconductor.

wherein the p-type semiconductor is obtained by oxidizing, with an oxidizing agent or through electrochemical doping, an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in claim 21, and


wherein the n-type semiconductor is obtained by reducing, with a reducing agent or through electrochemical doping, an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in claim 21.

33. **(Currently amended)** An organic electroluminescent device made by use of an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

34. **(Currently amended)** A non-linear organic material made by use of an aminoquinoxaline compound as defined in claim 1 or a polyaminoquinoxaline compound as defined in ~~claim 1~~ claim 21.

35. **(New)** A polyaminoquinoxaline compound having recurring units of the following formula (18a') obtained by polymerizing the monomer of claim 14,





wherein  $R^3$  and  $R^4$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y provided that when  $R^3$  and  $R^4$  are, respectively, the above-defined phenyl, pyridyl, biphenyl, naphthyl, thienyl, pyrrolyl, furyl or condensed heteroaryl group, these groups may be joined through a single bond;

$X^2$  represents  $-NH-R^5-NH_2$  or  $-NH-R^6$ ;

$R^5$  represents a  $C_1$ - $C_{10}$  alkylene group, a  $-C(O)CH_2-$ ,  $-CH_2C(O)-$ , a divalent benzene ring which may be substituted with Y, a divalent pyridine ring which may be substituted with Y, a divalent biphenyl group which may be substituted with Y, a divalent naphthalene ring which may be substituted with Y, a divalent thiophene ring which may be substituted with Y, a pyrrole

ring which may be substituted with Y, a furan ring which may be substituted with Y, or a condensed hetero ring which may be substituted with Y;

$R^6$  represents a hydrogen atom, a  $C_1$ - $C_{10}$  alkyl group, an acetyl group, a phenyl group which may be substituted with Y, a pyridyl group which may be substituted with Y, a biphenyl group which may be substituted with Y, a naphthyl group which may be substituted with Y, a thienyl group which may be substituted with Y, a pyrrolyl group which may be substituted with Y, a furyl group which may be substituted with Y or a condensed heteroaryl group which may be substituted with Y;

Y represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_1$ - $C_{10}$  cyanoalkyl group, a phenyl group which may be substituted with Z, a pyridyl group which may be substituted with Z, a biphenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z, a thienyl group which may be substituted with Z, a pyrrolyl group which may be substituted with Z, a furyl group which may be substituted with Z or a condensed heteroaryl group which may be substituted with Z provided that if Y is two or more in number, Y may be the same or different;

A<sup>2</sup> are each C or N,  $R^{59}$ - $R^{66}$  independently represent a hydrogen atom, a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group which may be substituted with Z, a naphthyl group which may be substituted with Z or a thienyl group which may be substituted with Z, provided that when A<sup>2</sup> represents N,  $R^{62}$  and  $R^{63}$  are both non-existent;

Z represents a halogen atom, a cyano group, a nitro group, an amino group, an epoxy group, a vinyl group, a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  haloalkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a

C<sub>1</sub>-C<sub>10</sub> cyanoalkyl group, a phenyl group, a biphenyl group, a naphthyl group, a thienyl group, a pyrolyl group, a furyl group or a condensed heteroaryl group, provided that if Z is two or more in number, Z may be the same or different; and

n is an integer of 2 or more.